



Jeopardy Game

Precalculus

Functions

Limits

Derivative

Evaluation of
derivatives

Theory

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Precalculus for 100.

$$\ln \frac{x}{y} =$$

$\ln x + \ln y$

$\ln x - \ln y$

$x \ln y$

$y \ln x$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Precalculus for 200.

The function $y = x^2 \cdot \sin x$ is

odd

even

neither odd nor even

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Precalculus for 300.

$\arctan 1 =$

∞
 π
 $-\frac{3\pi}{4}$
 $-\frac{4\pi}{6}$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Precalculus for 400.

The equivalence " $a < b$ if and only if $f(a) < f(b)$ " is the property of

- even functions
- one-to-one functions
- continuous functions
- increasing functions
- none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Functions for 100.

How many points of inflection is on the graph of the function $y = \sin x$ in the open interval $(0, 2\pi)$

- none
- one
- two
- three
- none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Functions for 200.

Find points of discontinuity of the function $y = \frac{x-4}{(x-2)\ln x}$

none

0

0, 1

0, 1, 2

0, 2

0, 1, 4

0, 4

none of them

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Functions for 300.

Let f be a function and f^{-1} be its inverse. Then $f^{-1}(f(x)) =$

0

1

x

$f(x)$

$f^{-1}(x)$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Functions for 400.

$\arcsin(\sin x) = x$ for every $x \in \mathbf{R}$

Yes

No

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Limits for 100.

$$\lim_{x \rightarrow -\infty} \operatorname{arctg} x =$$

0
 $\frac{\pi}{2}$
 $-\frac{\pi}{2}$

∞

$-\infty$

none of them

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Limits for 200.

$$\lim_{x \rightarrow \infty} \sin x =$$

1

-1

does not exist

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Limits for 300.

$$\lim_{x \rightarrow \infty} \frac{2x^3 + x^2 + 4}{x^2 - x + 2} =$$

∞

2

0

none of them

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Limits for 400.

$$\lim_{x \rightarrow 0^+} \frac{e^{1/x}(x - 1)}{x}$$

0

1

e

∞

-1

$-e$

$-\infty$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Derivative for 100.

$$\left(\frac{1}{\sqrt[3]{x}}\right)' =$$

$$\frac{1}{3}x^{-2/3}$$

$$-\frac{1}{3}x^{-2/3}$$

$$-\frac{1}{3}x^{1/3}$$

$$\frac{1}{3}x^{-4/3}$$

$$-\frac{1}{3}x^{-4/3}$$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Derivative for 200.

$$(x - x \ln x)' =$$

- $\ln x$
- $- \ln x$
- $1 + \ln x$
- $1 - \ln x$
- 0
- $1 - \frac{1}{x}$
- none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Derivative for 300.

$$(x^2 e^{x^2})'$$

$$2xe^{2x}$$

$$2xe^{x^2} 2x$$

$$2xe^{x^2} + x^2 e^{x^2}$$

$$2xe^{x^2} + x^2 e^{x^2} 2x$$

$$2xe^{x^2} 2x + x^2 e^{x^2} 2x$$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Derivative for 400.

The definition of the derivative of the function f at the point a is

$$\lim_{h \rightarrow 0} \frac{f(x + h) + f(x)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(x + h)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(x) - f(x + h)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(x - h) - f(x)}{h}$$

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Evaluation of derivatives for 100.

$$(x^2 + 1)' =$$

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Evaluation of derivatives for 200.

$$(xe^x)' =$$

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Evaluation of derivatives for 300.

$\ln(\sin x) =$

Robert Mařík
Jeopardy game

Home Page

GameBoard

Full Screen

Close

Quit



Evaluation of derivatives for 400.

$$(xe^{-x})' =$$

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Theory for 100.

By theorem of Bolzano, the polynomial $y = x^3 + 2x + 4$ has zero on

- (0, 1)
- (1, 2)
- (2, 3)
- (-1, 0)
- (-2, -1)
- (-3, -2)

none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Theory for 200.

Let $a \in Im(f)$. Then the solution of the equation $f(x) = a$ exists. This solution is unique if and only if

- f is one-to-one
- f is increasing
- f continuous
- f differentiable
- none of them

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Theory for 300.

If the function has a derivative at the point $x = a$, then it is

- increasing at a .
- decreasing at a .
- one-to-one at a .
- continuous at a .
- undefined at a .

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Theory for 400.

If both $y(a) = y'(a) = y''(a) = 0$, then the function

has local maximum at a .

has local minimum at a .

has point of inflection at a .

any of these possibilities may be true, we need more informations.

Robert Mařík
Jeopardy game

[Home Page](#)

[GameBoard](#)

[Full Screen](#)

[Close](#)

[Quit](#)